



Web Images Video News Maps more »

(array) and "common probe" and subfamily

2003

Search

Ad  
Sc  
Ss

The "AND" operator is unnecessary -- we include all search terms by default. [details]

Scholar All articles - Recent articles Results 1 - 10 of about 13 for (array) and "common probe" ar

### Long-term global gene expression patterns in irradiated human lymphocytes - all 6 versions »

S Fält, K Holmberg, B Lambert, A Wennberg - Carcinogenesis, 2003 - Oxford Univ Press  
... Small inducible cytokine subfamily C (probe id. ... cultures and clones were linked and common probe numbers were ... In a study using an array with approximately 6700 ...

Cited by 13 - Related Articles - Web Search - BL Direct

### Olfaction: Transient expression of a putative odorant receptor in the avian notochord - all 6 versions »

S Nef, P Nef - Proceedings of the National Academy of Sciences, 1997 - National Acad Sciences  
... family belongs to the large multigene superfamily of seven ... Chicken genomic DNA blot analysis of the COR7 subfamily. ... COR7b probes or with a common probe (COR7ab ...

Cited by 25 - Related Articles - Web Search - BL Direct

### Trypanosome glucose transporters - all 4 versions »

MF Barrett, E Tetaud, A Seyfang, F Bringaud, T ... - Molecular & Biochemical Parasitology, 1998 - Elsevier  
... These include: a conserved array of cysteine residues ... topology of the glucose transporter superfamily in that ... THT1 and THT2 genes (common probe), specific to ...

Cited by 23 - Related Articles - Web Search

### Identification of in vivo mRNA targets of GLD-1, a maxi-KH motif containing protein required for C. ... - all 10 versions »

MH Lee, T Schedl - Genes & Development, 2001 - Cold Spring Harbor Lab  
... binding protein of the GSG/STAR subfamily, controls diverse ... FLAG expression from animals rescued by complex arrays. ... Probe 11 also has a similar second sequence ...

Cited by 58 - Related Articles - Web Search - BL Direct

### Multiple Isoforms of Heparan Sulfate d-Glucosaminyl 3-O-Sulfotransferase ISOLATION, CHARACTERIZATION ... - all 3 versions »

NW Shworak, J Liu, LM Petros, L Zhang, M Kobayashi ... - Journal of Biological Chemistry, 1999 - ASBMB  
... side chains enables interactions with a broad array of protein ... regions reveals the human 3-OST-3 multigene subfamily. ... 3'B), and the 3-OST-3 common probe (ST-3 ...

Cited by 107 - Related Articles - Web Search - BL Direct

### The orphan receptor BmHNF-4 of the silkworm Bombyx mori: ovarian and zygotic expression of two mRNA ... - all 4 versions »

L Swavers, K Iatrou - Mechanisms of Development, 1998 - Elsevier  
... Because silkworm ovarioles consist of long arrays of more ... members of the nuclear hormone receptor superfamily (Fig. ... probes, 12 h for the 'common' probe and 4 ...

Cited by 11 - Related Articles - Web Search

### The orphan nuclear receptors BmE75A and BmE75C of the silkworm Bombyx mori: hormonal control and ... - all 4 versions »

L Swavers, T Eystathiou, K Iatrou - Insect Biochemistry and Molecular Biology, 2002 - Elsevier  
... simultaneously present and arranged in a linear array. ... for all BmE75 isoforms (common probe) are, generally ... encoding a nuclear receptor superfamily member: an ...